



agriculture

Department:
Agriculture
REPUBLIC OF SOUTH AFRICA

PHYTOSANITARY IMPORT REQUIREMENTS FOR FRESH *LYCOPERSICON* SPP. (TOMATO) FROM NAMIBIA TO SOUTH AFRICA

1. Additional Declaration on the Phytosanitary Certificate

1.1. The tomato in this consignment originates from NPPOZA's approved production site(s), packhouse(s), and storage facility(ies) that are registered by the NPPO of Namibia; received by the NPPOZA at least four weeks prior to the departure of the first consignment.

1.2. The tomato has been produced and packed according to the *Bactrocera dorsalis* risk mitigation measures as prescribed in Addendum 1.

OR

The tomato in this consignment originates in from production area (s), on the basis of the official annual surveys, *Bactrocera dorsalis* does not occur.

2. Registration of production sites, packhouses and storage facilities

2.1. *Lycopersicon* spp. for export to South Africa shall originate from production sites, packhouses and storage facilities that are approved and registered annually by the National Plant Protection Organisation (NPPO) of Namibia.

2.2. The list/database of the registered facilities that have been approved for export of *Lycopersicon* spp. to South Africa shall contain the following information:

2.2.1 Name and unique identification code of each production site, and the area in which the production site is situated.

2.2.2 Name and unique identification code of each pack house.

2.2.3 Name and unique identification code of each storage facility.

2.3. The list/database of the registered facilities that have been inspected, approved and registered by the NPPO of Namibia for the exportation of *Lycopersicon* spp. to South Africa shall be made available to the National

Plant Protection Organisation of South Africa (NPPOZA) annually. The NPPO of Namibia shall send the list of registered facilities to the NPPOZA at least four weeks prior to the departure of the first consignment. The NPPOZA shall assess the list/database and the approved facilities will be published on the Department of Agriculture (DoA) website.

- 2.4. The NPPO of Namibia shall ensure that *Lycopersicon* spp. for export to South Africa shall only originate from production area(s) which comply with these phytosanitary import requirements.

3. Pest free areas – *Bactrocera dorsalis*

- 3.1. The pest free status in respect to the fruit fly *Bactrocera dorsalis* in the *Lycopersicon* spp. production area(s) of Namibia will be accepted, in accordance with the guidelines outlined in ISPM 26: *Establishment of pest free areas for fruit flies (Tephritidae)*. (FAO, 2015).
- 3.2. The area from which *Lycopersicon* spp. is sourced (produced, handled and packed) for export to South Africa under *Bactrocera dorsalis* pest free area requirements shall have a detection survey system to verify the absence of this pest; and a pest monitoring system to verify that freedom from this pest has been maintained. Summary data including number and location of traps, data on trap catches, shall be made available to the NPPOZA upon request.
- 3.3. The NPPO of Namibia shall notify the NPPOZA within 4 business days when detections of *Bactrocera dorsalis* result in a change to the pest free areas.
- 3.4. The NPPO of Namibia shall work collaboratively with the NPPOZA to ensure that appropriate measures are implemented to mitigate the risk or that the pest free status is reinstated.

4. Post-harvest measures

- 4.1. Tomato shall be harvested in a hard condition with unbroken skin and that no tomato are harvested for export to South Africa showing evidence of premature ripening or yellowing.

- 4.2. Tomato destined for South Africa shall not be mixed with tomato destined for other markets in pack houses or storage facilities. *Lycopersicon* spp. shall be appropriately inspected, packed, stored and transported.
- 4.4. Rejected tomato shall be removed from the packing area at the end of each day.
- 4.5. Post-harvest inspections shall be conducted according to the ISPM 31: *Methodologies for sampling consignments* (FAO, 2008). This should be able to identify with at least 95% reliability; a level of infection of 0.5% or above.
- 4.6. Should any quarantine pest of concern (as listed in Annex 1) be detected, the consignment shall be rejected for export to South Africa. Namibia will lose its pest free status and have to inform the NPPO of South Africa immediately. The NPPO of South Africa will then have to advise the NPPO of Namibia of the phytosanitary action(s) to be taken, which can include the possibility of the temporary suspension of all exports until such time as an acceptable mitigation measure is agreed upon for future use.
- 4.7. The tomato in the consignment is free from leaves, plant debris and soil.
- 4.8. Only mature and symptomless tomato shall be packed for export to South Africa.
- 4.9. The packing materials for *Lycopersicon* spp. destined for South Africa shall be new and clean cardboard boxes/cartons or clean plastic crates or clean bulk bins.
- 4.10. All boxes or pallets shall be covered by insect proof netting or sheeting with holes not more than 1mm in diameter or length.
- 4.11. No packaging material of plant origin, including straw, shall be used.
- 4.12. Should wood packaging material be used, it shall comply with ISPM 15: *Regulation of wood packaging material in international trade* (FAO, 2009).

5. Marking requirements

- 5.1. Each carton (box) of *Lycopersicon* spp. tomato shall be marked in English with correct and accurate information.

6. Phytosanitary certification

- 6.1. An import permit is required in terms of the Agricultural Pests Act, 1983 (Act No. 36 of 1983) and associated Regulations R.111 of 27 January 1987 as

amended.

- 6.2. Upon completion of sampling and inspection, a Phytosanitary Certificate shall be issued by the NPPO of Namibia prior to shipment. Entry of the consignment to South Africa shall be subject to the availability of the original Phytosanitary Certificate. A Phytosanitary Certificate shall only be issued for *Lycopersicon* spp. that meet these phytosanitary requirements.

7. Phytosanitary inspection on arrival

- 7.1. Once a consignment of *Lycopersicon* spp. arrives at the designated port of entry, NPPOZA shall examine the relevant documents and markings.
- 7.2. Any consignment with certification that does not conform to specifications as set out in this phytosanitary import requirements shall be rejected.
- 7.3. Upon arrival of the consignment at the designated port of entry, a representative sample shall be drawn and inspected for quarantine pests and suspicious tomato shall be dissected to determine the status of infestation. Should pests or symptoms be found, the samples shall be sent for laboratory identification, and the consignment shall be detained pending the laboratory results.
- 7.4. If any pest(s) in Annex 1 are detected, Namibia will lose its pest free status. NPPOZA will immediately take the necessary corrective actions and notify the NPPO of Namibia. NPPOZA will advise the NPPO of Namibia of the action/s to be taken, which can include the possibility of the temporary suspension of all exports until such time as an acceptable mitigation measure is agreed upon before trade commences.
- 7.5. Should any potential quarantine pest that has not been categorized be detected on *Lycopersicon* spp. from Namibia, it shall require assessment to determine its quarantine status and whether phytosanitary action is required. The detection of any potential quarantine pest of concern not already identified in the analysis may result in a review of trade to ensure that phytosanitary measures provide the appropriate level of phytosanitary protection for South Africa.
- 7.6. The importer is responsible for all costs relating to disposal, removal or rerouting, including costs incurred by NPPOZA to monitor the action taken.

ANNEX 1: QUARANTINE PESTS OF CONCERN TO SOUTH AFRICA NOT OCCURRING ON *LYCOPERSICON* SPP. IN NAMIBIA

Viruses: Lettuce infectious yellows closterovirus
 Squash leaf curl bigeminivirus
 Squash mosaic comovirus
 Tobacco ringspot nepovirus
 Tomato ringspot nepovirus

Insects: *Anastrepha grandis* [Tephritidae]
Bactrocera cucumis [Tephritidae]
Bactrocera dorsalis (complex) [Tephritidae] Including: *B. carambolae*, *B. caryeae*, *B. kandiensis*, *B. occipitalis*, *B. pyrifoliae*
Bactrocera lycopersicone [Tephritidae]
Bactrocera latifrons [Tephritidae]
Bactrocera tau [Tephritidae]
Dacus demmerezi [Tephritidae]
Dacus solomonensis [Tephritidae]
Ceroplastes ceriferus [Coccidae]
Ceroplastes floridensis [Coccidae]
Maconellicoccus hirsutus [Pseudococcidae]
Rastrococcus iceryoides [Pseudococcidae]
Dysmicoccus lycopersicone [Pseudococcidae]
Dysmicoccus neobrevipes [Pseudococcidae]
Pseudococcus jackbeardsleyi [Pseudococcidae]
Diaphania nitidalis [Pyralidae]
Diaphana hyalinata [Pyralidae]
Thrips hawainensis [Thripidae]
Thrips palmi [Thripidae]
Zeugodacus cucurbitae [Tephritidae]

ADDENDUM 1: RISK MITIGATION MEASURES FOR *LYCOPERSICON* SPP. FROM NAMIBIA TO SOUTH AFRICA

The following pre- and post-harvest practices reflects the current system for risk management overseen by the NPPO of Namibia, employed by producers of *Lycopersicon* spp. to be imported to South Africa:

TABLE 1. OVERVIEW OF THE SYSTEM FOR THE COMMERCIAL PRODUCTION AND EXPORT OF *LYCOPERSICON* SPP. FROM NAMIBIA TO SOUTH AFRICA

ACTIVITIES	OUTCOMES
Pre-Harvest • In-field pest control activities • Good Agricultural Practice (GAP) e.g. cultural controls such as removal of weeds acting as pest reservoirs; pesticide application records; traceability system. • <i>Bactrocera dorsalis</i> control programme including a) seven day cycle field/orchard sanitation infestations b) application of insecticidal protein bait throughout the production cycle or Bait application technique (BAT) c) male annihilation throughout the production season with the placement of <i>Bactrocera dorsalis</i> respondent insecticidal male lures or the male annihilation technique(MAT)	• Reduced pre-harvest pest prevalence. • Reduced pre-harvest pest prevalence. • Reduced pest prevalence indicated by continuous monitoring of <i>Bactrocera dorsalis</i> to Male trap catch flies per trap per day value that is greater than 1 at any time during the production cycle may result in the suspension of the export program and the implementation of corrective measures.
Post-Harvest • Phytosanitary inspection	• Inspection of tomato and removal of external arthropod pests or

	infested/infected tomato or punctured/cracked tomato.
Regulatory/Official • Phytosanitary inspection and certification of consignments • Post-inspection product security • NPPOZA inspection of documentation and consignment on arrival in South Africa • Non-conformance contingencies • Pathway monitoring • Packing, labelling and storage compliance	• Certification by the NPPO of Namibia that consignments are free from regulated pests. • Prevention of post-treatment infestation of consignments by regulated pests e.g. pest-proof packaging. • Verification that the phytosanitary import requirements has been met. • Treat/re-ship/destroy non-conforming consignment. • Assurance that phytosanitary import requirements are being met. • Traceability and pest contamination

PRE-HARVEST ACTIVITIES

a) In-field pest control practices

- Namibian *Lycopersicon* spp. growers shall utilize pest control measures to reduce pre-harvest pest prevalence in commercially produced *Lycopersicon* spp. for export to other countries.
- These measures include a *Bactrocera dorsalis* control programme, and compliance with Good Agricultural Practice (GAP) as outlined below.

b) Good Agricultural Practice (GAP)

- The GlobalGAP standard for *Lycopersicon* spp. production requires training programmes for farmers and provincial government representative's safe use of agrichemicals, on-farm recording of fertilizer applications and crop protection products, inventory, sales, keeping receipts of input purchases and sales record-keeping, and safe tomato handling.
- The cultural control practices to be undertaken such as removal/suppression of weeds and fallen tomato, which act as reservoirs for pests.
- GAP is not a phytosanitary requirement but is advantageous for record keeping, particularly with respect to crop protection practices. GAP is also important for defining harvesting and post-harvest handling activities, traceability and recall throughout the export chain.

c) *Bactrocera dorsalis* control programme

- A specific programme shall be in place for *Bactrocera dorsalis* in Namibia should include surveillance to detect and determine species composition, and infestation rates.
- The programme shall be maintained by the NPPO of Namibia throughout the year in *Lycopersicon* spp. production areas.
- The surveillance programme shall incorporate trapping using cue lure and methyl eugenol, *Lycopersicon* spp. orchard surveys, periodic random and targeted cutting of tomato collected from orchards and local markets.
- A protein bait spray and insecticide shall be applied in the production areas for *Bactrocera dorsalis* control (Table 1).

POST-HARVEST ACTIVITIES

a) Pre-treatment procedures

Harvested *Lycopersicon* spp. shall be:

- Covered with insect proof material to prevent re-infestation by arthropods during transit to primary pack houses or directly to packing facilities;
- Leaves and stems (but not bracts) are removed from the tomato;
- Damaged/infested/infected tomato is removed;

The above activities provide opportunity for operator inspection of tomato. Removal of damaged tomato can reduce the incidence of storage rots in tomato.

1. RISK MANAGEMENT MEASURES AND PHYTOSANITARY PROCEDURES (Table 1)

1.1. Management damaged tomato's/ infested tomato by external feeders

- Tomato with punctures/cracks or tomato damaged by external or surface-feeding arthropods shall not be packed for export to South Africa.

1.2. Management of *Bactrocera dorsalis*:

- The production site control program for *B. dorsalis* shall include an Integrated Pest Management (IPM) program using appropriate, effective and compatible measures at critical stages of development of the pest and crop.
- Population monitoring shall be based on production site inspections and forecasts of infestations.
- Information pertaining to production site control program for *B. dorsalis* shall be made available to NPPOZA on request (Table 1).

1.3. Supporting operational maintenance systems and verification of phytosanitary status

- A system of operational procedures shall be in place to ensure that the phytosanitary status of *Lycopersicon* spp. from Namibia is maintained and verified during the process of production and export to South Africa.
- The proposed system of operational maintenance for the production and export of *Lycopersicon* spp. from Namibia to South Africa consists of:
 - ✓ pre-export inspection by the NPPO of Namibia;
 - ✓ packaging and labelling compliance;
 - ✓ phytosanitary certification by NPPO of Namibia;
 - ✓ specific conditions for storage and movement; and
 - ✓ on-arrival quarantine inspection by NPPOZA in South Africa.

A. Pre-export inspection and remedial action by the NPPO of Namibia

- ✓ The NPPO of Namibia shall inspect all consignments in accordance with official procedures for all quarantine pests using sampling procedures developed by NPPOZA as outlined.
- ✓ If actionable pests as listed are found during these inspections, then remedial action shall be taken.
- ✓ Records of the interceptions made during these inspections (live or dead quarantine pests) shall be maintained by the NPPO of Namibia and made available to NPPOZA if requested.
- ✓ If *Bactrocera dorsalis* is detected, the consignment shall be rejected for export to South Africa.

B. Packing and labelling

- ✓ All packages of *Lycopersicon* spp. for export shall be free from contaminated plant materials including soil, splinters, twigs, leaves and other plant materials.
- ✓ Inspected and treated *Lycopersicon* spp. shall be packed in new boxes.
- ✓ No unprocessed packing material of plant origin, such as straw, shall be allowed.
- ✓ All wood material used in packaging of *Lycopersicon* spp. shall comply with ISPM 15.
- ✓ All boxes shall be labelled with the name of production site and name pack house for the purposes of trace back if necessary.

C. Phytosanitary certification by the NPPO of Namibia

- ✓ Before a phytosanitary certificate is issued, the NPPO of Namibia shall conduct phytosanitary inspection to ensure that the number of packaged tomato is consistent with the number of disinfested tomato's, traceability labelling is complete, packaging is insect-proof, the tomato is free from regulated pests and that all other importing country requirements have been met.
- ✓ The NPPO of Namibia shall issue a Phytosanitary Certificate for each consignment upon completion of pre-export phytosanitary inspection.
- ✓ Each Phytosanitary Certificate is to contain the following information:

Distinguishing marks

- ✓ The names of production site and pack house, together with the number of boxes per consignment. This is to ensure trace back to the production site in the event that this is necessary.

D. Specific conditions for storage and movement

- ✓ Packed product and packaging shall be protected from pest contamination during and after packing, during storage and during movement between locations (e.g. pack house to cool storage/depot, to inspection point, to export point).
- ✓ *Lycopersicon* spp. for export to South Africa shall be inspected and certified by the NPPO of Namibia, and shall be maintained in secure conditions to prevent mixing with *Lycopersicon* spp. for export to other destinations or the domestic market and kept in secure storage until export.

E. On-arrival quarantine inspection and remedial action, and clearance by NPPOZA in South Africa

- ✓ On arrival in South Africa, each consignment shall be inspected by NPPOZA.
- ✓ *Lycopersicon* spp. from each consignment shall be randomly sampled for inspection. Such sampling methodology will provide for a 95% confidence level of detecting packing units with infested *Lycopersicon* spp. if the infestation rate is 2% or higher.
- ✓ If actionable quarantine pests are found during these inspections, then remedial action shall be taken.
- ✓ Where consignments are found to be non-compliant with requirements on-arrival, the importer shall be given the option to treat (if suitable treatments for the pests detected can be applied), re-export or destroy the consignment.

ADDENDUM 2: NATIONAL QUARANTINE PESTS LIST FOR LYCOPERSICON SPP. (TOMATO) FOR SOUTH AFRICA

Viruses: Lettuce infectious yellows closterovirus
 Squash leaf curl bigeminivirus
 Squash mosaic comovirus
 Tobacco ringspot nepovirus
 Tomato ringspot nepovirus

Insects: *Anastrepha grandis* [Tephritidae]
Bactrocera cucumis [Tephritidae]
Bactrocera lycopersicone [Tephritidae]
Bactrocera dorsalis [Tephritidae]
Bactrocera dorsalis (complex) [Tephritidae] Including: *B. carambolae*, *B. caryeae*, *B. kandiensis*, *B. occipitalis*, *B. pyrifoliae*
Bactrocera latifrons [Tephritidae]
Bactrocera tau [Tephritidae]
Dacus demmerezi [Tephritidae]
Dacus solomonensis [Tephritidae]
Ceroplastes ceriferus [Coccidae]
Ceroplastes floridensis [Coccidae]
Maconellicoccus hirsutus [Pseudococcidae]
Rastrococcus iceryoides [Pseudococcidae]
Dysmicoccus Lycopersicone [Pseudococcidae]
Dysmicoccus neobrevipes [Pseudococcidae]
Prodiplosis longifila [Cecidomyiidae]
Pseudococcus jackbeardsleyi [Pseudococcidae]
Diaphania nitidalis [Pyralidae]
Diaphana hyalinata [Pyralidae]
Thrips hawainensis [Thripidae]
Thrips palmi [Thripidae]
Zeugodacus cucurbitae [Tephritidae]